

Structure genetique des flets (Platichthys flesus , Teleostei, Pleuronectidae)



Six populations of flounder (*Platichthys flesus*) from the Mediterranean and Atlantic coasts were analysed by enzymatic electrophoresis at 21 loci. The results confirm the separation of the species in three genetic groups. Differences between these genetic groups have also been observed.

Auteurs du document : Berrebi, P, Agnese, J, Vianet, R

Obtenir le document : Actes de colloques. Ifremer. Brest [ACTES COLLOQ. IFREMER.]. 1985

Mots clés : MED, France, A, Atlantic, *Platichthys flesus*, Pleuronectidae, Pisces, Taxonomy, Geographical distribution, Enzymes, Genes, Genotypes, Population genetics

Thème (issu du Text Mining) : MILIEU NATUREL, FAUNE, SCIENCES EXACTES SCIENCES HUMAINES

Date : 1983-12

Format : text/xml

Langue : Inconnu

Droits d'utilisation : info:eu-repo/semantics/openAccess, restricted use

Télécharger les documents : <https://archimer.ifremer.fr/doc/1983/acte-1242.pdf>

<https://archimer.ifremer.fr/doc/00000/1242/>

Permalien : <https://www.documentation.eauetbiodiversite.fr/notice/structure-genetique-des-flets-platichthys-flesus-teleostei-pleuronectidae0>

Evaluer cette notice: