

Etude de la composition corporelle de la sole (Solea solea) au cours du jeune: Influence de la température



The effects of starvation on Solea solea at 10 and 20 degree C in sea water, were studied. At 20 degree C fishes start to die on week 4; at 10 degree C they support a longer starvation period (9 weeks). Proximate analysis shows that protein are more steadily used at 20 degree C than at 10 degree C, contrarily to the corporal lipids which are used at the same rate. Starvation results in a reduction of liver lipids and glycogen. This reduction is more pronounced at 20 degree C : 70% of hepatic glycogen is used at this temperature within the first week of starvation against 30% at 10 degree C.

Auteurs du document : Richard, P, Djabali, M, Lagardere, Francoise, Lagardere, J

Obtenir le document : Actes de colloques. Ifremer. Brest [ACTES COLLOQ. IFREMER.]. 1985

Mots clés : Solea solea, Soleidae, Pisces, Survival, Tissues, Biochemical composition, Temperature effects, Proteins, Lipids, Starvation, Metabolism

Thème (issu du Text Mining) : FAUNE, AGRICULTURE, BIOCHIMIE - CHIMIE

Date : 1983-12

Format : text/xml

Langue : Inconnu

Droits d'utilisation : info:eu-repo/semantics/openAccess, restricted use

Télécharger les documents : <https://archimer.ifremer.fr/doc/1983/acte-1246.pdf>

<https://archimer.ifremer.fr/doc/00000/1246/>

Permalink : <https://www.documentation.eauetbiodiversite.fr/notice/etude-de-la-composition-corporelle-de-la-sole-solea-solea-au-cours-du-jeune-influence-de-la-temperat0>

[Evaluer cette notice:](#)