

Nouvelles especes de Nannastacidae (Crustaces, Cumaces) dans les eaux sahariennes de l'Atlantique.



While only 4 species were known for the family Nannastacidae in West Africa, a recent study has revealed 11 more species - *Procampylaspis maurini*, *P. arguini*, *P. thalassae*, *Cumella africana*, *Schizotrema atlantica*, *Campylaspis macrophthalma*, *C. mauritanica*, *C. bonetti*, *C. legendrei africana*, *C. jonesi*. Of these, *S. atlantica* and the last 4 *Campylaspis* species are new species, and are described. Both male and female adults are described, with explanatory diagrams, and additional observations are included. The species of *Campylaspis* are incorporated modified form. It is thought that this description completes not only acquaintance with the fauna of the Saharian section of the Atlantic, but also the whole of the Atlantic Ocean.

Auteurs du document : Bacescu, M., Muradian, Zarui

Obtenir le document : ISTPM

Mots clés : Species, Saharian waters, Nannastacidae Crustaceans, Indexing in process

Thème (issu du Text Mining) : MILIEU NATUREL, INDUSTRIE, FAUNE, INFORMATION - INFORMATIQUE

Date : 1972-03

Format : text/xml

Source : Revue des Travaux de l'Institut des Pêches Maritimes (0035-2276) (ISTPM), 1972-03 , Vol. 36 , N. 3 , P. 255-269

Langue : Inconnu

Droits d'utilisation : Ifremer, info:eu-repo/semantics/openAccess, restricted use

Télécharger les documents : <https://archimer.ifremer.fr/doc/1972/publication-2046.pdf>

<https://archimer.ifremer.fr/doc/00000/2046/>

Permalien : <https://www.documentation.eauetbiodiversite.fr/notice/nouvelles-especes-de-nannastacidae-crustaces-cumaces-dans-les-eaux-sahariennes-de-l-atlantique0>