

Consequences du cycle de mue et d'intermue sur la morphologie fonctionnelle des pars molaris des Crustacés Peracarides.



The anatomy of the masticatory part of the mandible (the pars molaris) has been studied throughout the moult/intermoult cycle in some peracarid crustacean species, by scanning electron microscopy. Species examined were: *Ligia oceanica*, *Sphaeroma serratum*, *Idotea balthica*, *Talitrus saltator*, *Hyale nilssoni*, Gammarid amphipoda and the mysid *Pranus flexuosus*. Feeding is known to decline near moulting; this may be because the chewing surfaces become loosened before moulting. Also the surfaces are worn down and useless, and feeding ceases. Very long intermoult periods, as found in old individuals, may thus cause long periods of starvation and this could enhance the symptoms of senescence.

Auteurs du document : Saudray, Yves

Obtenir le document : ISTPM

Mots clés : Peracarid crustaceans, Morphology, Consequences, Indexing in process

Thème (issu du Text Mining) : FAUNE, INFORMATION - INFORMATIQUE

Date : 1972-03

Format : text/xml

Source : Revue des Travaux de l'Institut des Pêches Maritimes (0035-2276) (ISTPM), 1972-03 , Vol. 36 , N. 3 , P. 273. 1972

Langue : Inconnu

Droits d'utilisation : Ifremer, info:eu-repo/semantics/openAccess, restricted use

Télécharger les documents : <https://archimer.ifremer.fr/doc/1972/publication-2047.pdf>

<https://archimer.ifremer.fr/doc/00000/2047/>

Permalien : <https://www.documentation.eauetbiodiversite.fr/notice/consequences-du-cycle-de-mue-et-d-intermue-sur-la-morphologie-fonctionnelle-des-pars-molaris-des-cru0>

Evaluer cette notice: