

La reproduction de l'Ormeau *Haliotis tuberculata* L.



In order to improve the regulation of abalone 'fishing' and to provide for the restocking of supplies, more details of the reproductive cycle were sought. Attention is given to gonad morphology, acquisition of sexual maturity, juvenile hermaphroditism, gonad histogenesis and histology of gametogenesis, annual cycle of gamete emission, and the relation between size and fertility. Results are presented in graphs, histograms, diagrams and photographs. It was found that female reached sexual maturity during the 3rd year, at 38-54 mm, male during year 2 at 25-40 mm; there was successive juvenile hermaphroditism; an active period from mid-June to mid-November was indicated by the volumetric index of maturity; water temperature was associated with the reproductive cycle, the highest temperatures coinciding with maximum for 7.5 cm individuals increasing to $> 5 \times 10$ super(6) for those of 11.5 cm.

Auteurs du document : Girard, Annie

Obtenir le document : ISTPM

Mots clés : Ormeau, Cycle, Temperature, *Haliotis tuberculata* L., Reproduction

Thème (issu du Text Mining) : MILIEU NATUREL, SCIENCES EXACTES SCIENCES HUMAINES, INFORMATION - INFORMATIQUE

Date : 1972-06

Format : text/xml

Source : Revue des Travaux de l'Institut des Pêches Maritimes (0035-2276) (ISTPM), 1972-06 , Vol. 36 , N. 2 , P. 163-184

Langue : Inconnu

Droits d'utilisation : Ifremer, info:eu-repo/semantics/openAccess, restricted use

Télécharger les documents : <https://archimer.ifremer.fr/doc/1972/publication-2062.pdf>

<https://archimer.ifremer.fr/doc/00000/2062/>

Permalien : <https://www.documentation.eauetbiodiversite.fr/notice/la-reproduction-de-l-ormeau-haliotis-tuberculata-10>