

La diatomee *Navicula ostrearia* Bory en baie de Bourgneuf



Previously, *Navicula ostrearia* B. has been described only from the Marennes-Oleron district of the Atlantic coast of France. However, this pennate diatom is also present in the Bay of Bourgneuf where breeding of green-gilled oysters were recognized in sea basins called oyster ponds. A short description of observations made with the optical microscope precedes description of high magnification examination of the frustule made with the scanning electron microscope. This combination of techniques shows a polymorphism between the external and internal surfaces of diatom valves.

Auteurs du document : Robert, Jean-michel

Obtenir le document : ISTPM

Mots clés : North east Atlantic, France, Oyster pond, Microscopic observation, Polymorphism, Diatom

Thème (issu du Text Mining) : MILIEU NATUREL, INFORMATION - INFORMATIQUE

Date : 1973-09

Format : text/xml

Source : Revue des Travaux de l'Institut des Pêches Maritimes (0035-2276) (ISTPM), 1973-09, Vol. 37, N. 3, P. 363-368

Langue : Inconnu

Droits d'utilisation : Ifremer, info:eu-repo/semantics/openAccess, restricted use

Télécharger les documents : <https://archimer.ifremer.fr/doc/1973/publication-2148.pdf>

<https://archimer.ifremer.fr/doc/00000/2148/>

Permalien : <https://www.documentation.eauetbiodiversite.fr/notice/la-diatomee-navicula-ostrearia-bory-en-baie-de-bourgneuf0>

Evaluer cette notice: