

Le "fleuve cotier": Un phenomene hydrologique important en Manche orientale. Exemple du Pas-de-Calais



The English Channel has a "megatidal" regime: in the Dover Strait, the tidal range varies between 3 and 9 m in neap tide and spring-tide respectively. The tide currents are alternating and essentially parallel to the coast. Along the French coast, the fluvial supplies, distributed from the Bay of Seine to Cape Gris-Nez, generate a coastal water mass which drifts nearshore, separated from the open sea by a frontal area: this coastal flow is characterized by its freshness, turbidity, phytoplankton richness, and pollution content. Its width is only 3 to 5 miles, but because of the high concentrations of dissolved and suspended components, it is possible that the coastal flow plays an important role in the fluxes between the Eastern Channel and the North Sea.

Auteurs du document : Brylinski, J M, Lagadeuc, Y, Gentilhomme, V, Dupont, J P, Lafite, R, Dupeuble, P A, Huault, M F, Auger, Y

Obtenir le document : Gauthier-Villars

Thème (issu du Text Mining) : MILIEU NATUREL

Date : 1991

Format : text/xml

Source : Oceanologica Acta, Special issue (0399-1784) (Gauthier-Villars), 1991

Langue : Inconnu

Droits d'utilisation : info:eu-repo/semantics/openAccess, restricted use

Télécharger les documents : <https://archimer.ifremer.fr/doc/00268/37874/35955.pdf>

<https://archimer.ifremer.fr/doc/00268/37874/>

Permalien : <https://www.documentation.eauetbiodiversite.fr/notice/le-fleuve-cotier-un-phenomene-hydrologique-important-en-manche-orientale-exemple-du-pas-de-calais0>

[Evaluer cette notice:](#)