


## Leydig cells in



Males of *Myleus ternetzi*, at various maturation stages were netted in a French Guyana river. Testes were histologically and ultrastructurally analysed. Leydig cells were always present in large numbers. In mature or almost mature testes, they were characterized by a large spherical nucleus and an extensively developed smooth endoplasmic reticulum (SER) composed of very narrow tightly packed tubules. In regressed testes, most of the Leydig cells had a smaller nucleus with an irregular shape, a disorganized SER and the fine structure of the mitochondria was greatly affected. These changes would reflect different phases throughout a metabolic cycle in relation to spermatogenesis.

**Auteurs du document :** Maurice Loir, Anne Margeridon, Chantal Cauty

**Obtenir le document :** EDP Sciences

**Mots clés :** Fish, South America, testis, Leydig cell, ultrastructure, Poisson, Amérique du Sud, testicule, cellule de

Leydig, ultrastructure

**Date :** 1989-01-15

**Format :** text/xml

**Source :** <https://doi.org/10.1051/alr:1989006>

**Langue :** Anglais

**Télécharger les documents :** <https://www.alr-journal.org/10.1051/alr:1989006/pdf>

**Permalien :** <https://www.documentation.eauetbiodiversite.fr/notice/leydig-cells-in0>

[Evaluer cette notice:](#)



Ce portail, créé et géré par l'Office International de l'Eau (OIEau), est géré avec l'appui de l'Office français de la biodiversité (OFB)

