

## Diet of invasive pikeperch



Impact assessments of invasive piscivorous fishes usually rely on dietary analyses to quantify their predation pressure on prey communities. Stomach contents analysis (SCA), typically a destructive sampling method, is frequently used for this. However, many invasive piscivores are exploited by catch-and-release sport angling, with destructive sampling often not feasible. Stable isotope analysis (SIA) provides an alternative dietary analysis tool to SCA, with use of fin tissue, scales and/or epidermal mucus potentially enabling its non-destructive application. Here, the diet of a population of pikeperch *Sander lucioperca*, an invasive sport fish to Great Britain, was investigated by applying SIA to a range of tissues. Testing SI data of dorsal muscle (destructive sampling) versus fin, scale and mucus (non-destructive sampling) revealed highly significant relationships, indicating that the tissues collected non-destructively can be reliably applied to pikeperch diet assessments. Application of these SI data to Bayesian mixing models predicted that as *S. lucioperca* length increased, their diet shifted from macro-invertebrates to fish. Although similar ontogenetic patterns were evident in SCA, this was inhibited by 54% of fish having empty stomachs. Nevertheless, SCA revealed that as *S. lucioperca* length increased, their prey size significantly increased. However, the prey:predator length ratios ranged between 0.08 and 0.38, indicating most prey were relatively small. These results suggest that when non-destructive sampling is required for dietary analyses of sport fishes, SIA can be applied using fin, scales and/or mucus. However, where destructive sampling has been completed, SCA provides complementary dietary insights, especially in relation to prey size.

**Auteurs du document** : Emma T. Nolan, J. Robert Britton

**Obtenir le document** : EDP Sciences

**Mots clés** : Bayesian mixing models, gut contents, piscivory, trophic, modèles bayésiens de mélange, carbone, contenu intestinal, azote, piscivorie, écologie trophique

**Thème (issu du Text Mining)** : FAUNE, BIOCHIMIE - CHIMIE

**Date** : 2018-12-17

**Format** : text/xml

**Source** : <https://doi.org/10.1051/kmae/2018037>

**Langue** : Anglais

**Télécharger les documents** : <https://www.kmae-journal.org/10.1051/kmae/2018037/pdf>

**Permalien** : <https://www.documentation.eauetbiodiversite.fr/notice/diet-of-invasive-pikeperch0>

Evaluer cette notice: